

Tilburg University

Homomorphisms of graphs to odd cycles

Gerards, A.M.H.

Publication date:
1985

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Gerards, A. M. H. (1985). *Homomorphisms of graphs to odd cycles*. (Research Memorandum FEW). Faculteit der Economische Wetenschappen.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


17
R
CBM
R
7626
1985
185



* C I N O O 4 3 9 *

subfaculteit der econometrie

RESEARCH MEMORANDUM



TILBURG UNIVERSITY
DEPARTMENT OF ECONOMICS

Postbus 90153 - 5000 LE Tilburg
Netherlands



FEW
185

5077

1977

HOMOMORPHISMS OF GRAPHS TO ODD CYCLES

A.H.M. GERARDS

HOMOMORPHISMS OF GRAPHS TO ODD CYCLES

A.M.H. GERARDS^{*})

ABSTRACT. We give a class of graphs G for which there exists a homomorphic (= adjacency preserving) map from $V(G)$ to $V(C)$, where C is the shortest odd cycle in G . Hereby extending a result of Albertson, Catlin and Gibbons. Moreover our class of graphs is characterized by the property: For each odd subdivision of G' there exists a homomorphic map from $V(G')$ to $V(C')$ where C' is the shortest odd cycle of G' .

^{*}) Department of Econometrics
Tilburg University, P.O. Box 90153, Tilburg, Holland.

1. INTRODUCTION

A graph G' is called a homomorphic image of a graph G if there exists a map $\varphi : V(G) \rightarrow V(G')$ preserving adjacency, i.e. if $uv \in E(G)$ then $\varphi(u)\varphi(v) \in E(G')$ ($V(G)$ denotes the nodeset of G , $E(G)$ its edgeset). We say G maps to G' . In this paper we are interested in maps to odd cycles. Obviously, if G maps to an odd cycle of length M then no odd cycle in G is shorter than M . The result of this paper is

THEOREM. *Let G be a non-bipartite graph. Then either G maps to its shortest odd cycle or G contains an $\text{odd-}K_4$ or an $\text{odd-}K_3^2$ subgraph.* \square

Here an $\text{odd-}K_4$ and an $\text{odd-}K_3^2$ are graphs as indicated in fig. 1. Wiggled and dotted lines stand for (pairwise openly disjoint) paths, dotted lines may have length zero and *odd* indicates that the corresponding faces are odd cycles.



fig. 1

We shall give a proof in section 2. As pointed out in the remark at the end of that section the proof yields a polynomial-time algorithm which finds a map to the shortest odd cycle of G or finds an $\text{odd-}K_4$ or an $\text{odd-}K_3^2$. The theorem implies the following result due to Albertson, Catlin and Gibbons [1984]. Here *fold* means repeated identification of nodes at distance two.

COROLLARY 1. *Let G be a graph, and let M be an odd number (≥ 3) such that G contains no odd cycle shorter than M . Then either G maps to a cycle of length M or G contains a subgraph which folds to an $\text{odd-}K_4$ in which all odd cycles have length M .*

PROOF (that corollary 1 follows from the theorem).

If G is bipartite it maps to each odd cycle. So assume G is not bipartite. Let K be the length of its shortest odd cycle. Assume G does not map to a cycle of length M . Since each odd cycle of length greater than or equal to M maps to a cycle of length M , G does not map to a cycle of length K . By the theorem G has a subgraph G' such that G is an $\text{odd-}K_4$ or an $\text{odd-}K_3^2$. It is not hard to see that G' folds to a $\text{odd-}K_4$ in which each odd cycle has length M . \square

Note that the theorem properly extends corollary 1 since the graph in fig. 2 folds to K_4 (the 4-clique), but it does not contain an $\text{odd-}K_4$ or an $\text{odd-}K_3^2$.




fig. 2

As another corollary of the theorem we get a characterization of the graphs not containing an odd- K_4 or an odd- K_3^2 . The proof is left to the reader as an easy exercise. In [Gerards, Schrijver, 1985b] one can find more characterizations of these graphs. Using one of these characterizations Schrijver obtained a short and elegant proof of the main theorem of this paper. However that characterization relies on Tutte's characterization of regular matroids (Tutte [1958]), whereas the proof given in section 2 is elementary.

Odd-subdivision means: replacing edges by paths of odd length.

COROLLARY 2. *A graph G does not contain an odd- K_4 or an odd- K_3^2 iff G is bipartite or each odd-subdivision G' of G maps to the shortest odd cycle of G' .* □

REMARK. From the theorem it follows: If G contains no odd- K_4 and no odd- K_3^2 then the length of a shortest odd cycle equals the maximal cardinality of a packing of $E(G)$ with sets of the form $\langle X \rangle \cup \langle V(G) \setminus X \rangle$, $X \subset V(G)$. Here $\langle X \rangle$ denotes the set of edges with both ends in X . A weighted version of this min-max relations follows via odd-subdivision if all weights are odd. In fact Seymour [1977] showed that this weighted form holds for any weight function and for any signed graph not reducible to the signed graph in fig. 3. For signed graphs and reductions of them see the first part of section 2.

2. PROOF OF THE THEOREM

Before we get to the actual proof we give a preliminary result. It is convenient for its proof to state it in terms of signed graphs.

A *signed graph* (G, E_0) is an undirected graph G together with a subset, E_0 , of the edgeset $E(G)$. The edges in E_0 are called *odd*, the other edges are called *even*. A cycle C in G is called *odd (even)* if $E_0 \cap E(C)$ is *odd (even)* respectively). A signed graph is *bipartite* if it contains no odd cycles. Obviously odd- K_4 and odd- K_3^2 can be defined in this context as well. (Then the faces in fig. 1 indicated with *odd* are signed odd cycles.)

The following proposition is easy to check.

PROPOSITION. *A signed graph (G, E_0) contains an odd- K_4 or an odd- K_3^2 iff G can be reduced to K_4 (with all edges odd) or to the graph in fig. 3 (bold edges even, thin edges odd).* □

Reduction means: deletion of edges, contraction of even edges, or *resigning*, i.e. replacing E_0 by $E_0 \Delta B$, where B is a minimal cut ($\Delta :=$ "symmetric difference").



fig. 3

REMARK. Not that also holds " (G, E_0) contains an odd- K_4 iff (G, E_0) can be reduced to K_4 ". But " (G, E_0) contains an odd- K_3^2 iff (G, E_0) can be reduced to the signed graph in fig. 3" is not true.

We call a cycle C in a (signed) graph *non-separating* if for each two edges e and f not in $E(C)$ there exists nodes v_1, \dots, v_k not on C such that v_1 is contained in e , v_k is contained in f , and v_j and v_{j+1} are adjacent ($j = 1, \dots, k-1$). So C is separating if removing C from G (including the nodes of C) topologically disconnects G .

LEMMA. Let (G, E_0) be a signed graph not containing an odd- K_4 or an odd- K_3^2 . Let C be a non-separating odd cycle in (G, E_0) . If the edges induced by the nodes not on C form a bipartite signed graph then all odd cycles in G contain one fixed node of G .

PROOF.

If $V(G) \setminus V(C) = \emptyset$, then each edge in $E(G) \setminus E(C)$ is a chord of C . Since C is non-separating, $|E(G) \setminus E(C)| \leq 1$ in which case the lemma follows trivially. So let us assume: $V(G) \setminus V(C) \neq \emptyset$. Let T be a tree in G with $V(T) = V(G) \setminus V(C)$ (T exists since C is non-separating). Delete the edges induced by $V(G) \setminus V(C)$ not on T . Resign such that $E(T) \cap E_0 = \emptyset$. Contract the edges in $E(T)$. Now

each odd cycle in G contains an odd cycle in the contracted graph, which by the proposition does not contain an odd- K_4 or an odd- K_3^2 . Hence we may assume G to be the contracted graph. So $V(G) = V(C) \cup \{w\}$ (w not in $V(C)$). Let C' be an odd cycle in G , such that $|E(C') \cap E(C)|$ is minimal. Choose $u \in V(C) \cap V(C')$. We show that each odd cycle in G contains u . Suppose to the contrary that odd cycle C'' does not contain u . By the minimality of $|E(C') \cap E(C)|$ we have that $E(C') \cap E(C) \cap E(C'') = \emptyset$. There are five possibilities now, indicated in fig. 4. In any of them G contains an odd- K_4 or an odd- K_3^2 .




fig. 4

□

Now we leave the context of signed graphs. If we refer to the previous part of this section, we do this by referring to "the proposition" or "the lemma", and we assume $E_0 = E(G)$.

PROOF OF THE THEOREM. Let G be a minimal counterexample. Let M be the length of its shortest odd cycle. \mathcal{S} denotes the collection of shortest odd cycles in G . It is not hard to see that in a minimal counterexample, as G is, there cannot be a separating shortest odd cycle. Let $E_{\mathcal{S}}$ denote the set of edges contained in a shortest odd cycle. Let $G_1 = (V_1, E_1)$, $G_2 = (V_2, E_2)$, ..., $G_k = (V_k, E_k)$ be the components of $(V(G), E_{\mathcal{S}})$. (Note that (V_i, E_i) may be equal to $(\{v\}, \emptyset)$ for a $v \in V(G)$.)

Claim 1. Let $i = 1, \dots, k$. If Γ_1 and Γ_2 are odd cycles in G such that $V(\Gamma_1) \cap V_i \neq \emptyset$ and $V(\Gamma_2) \cap V_i \neq \emptyset$ then $V(\Gamma_1) \cap V(\Gamma_2) \neq \emptyset$,

Proof of claim 1. Assume Γ_1 and Γ_2 are odd cycles, $V(\Gamma_1) \cap V_i \neq \emptyset$, $V(\Gamma_2 \cap V_i) \neq \emptyset$ and $V(\Gamma_1) \cap V(\Gamma_2) = \emptyset$.

Since G_i is connected there exist C_1, C_2, \dots, C_ℓ in G_i , $C_j \in S$ ($j = 1, \dots, \ell$) such that $V(\Gamma_1) \cap V(C_1) \neq \emptyset$, $V(C_j) \cap V(C_{j+1}) \neq \emptyset$ ($j = 1, \dots, \ell-1$), and $V(\Gamma_2) \cap V(C_\ell) \neq \emptyset$. Without loss of generality we may assume: $\ell=1$. Let F be the set of edges in $E(\Gamma_1) \cup E(\Gamma_2)$ which do not meet C_1 . Then F is a forest. Extend F to a tree T such that $V(T) = V(G) \setminus V(C_1)$ (that is possible as C_1 is non-separating). Then the lemma applies to C_1 as a subgraph of $E(T) \cup E(\Gamma_1) \cup E(\Gamma_2) \cup E(C_1)$. Hence $V(\Gamma_1) \cap V(\Gamma_2) = \emptyset$, contradicting our assumption.

end of proof of claim 1

We are now going to prove that $k=1$, i.e. (V, E_G) is connected. We do this by contradiction.

Assumption: $k \geq 2$.

Let B be the set of edges leaving V_1 . Since G is connected, $B \neq \emptyset$. Moreover $e \in B$ implies that there is no $C \in S$ with $e \in E(C)$.

Claim 2. If Γ is an odd cycle in G , then $|E(\Gamma) \setminus B| \geq M$.

Proof of claim 2. Let Γ be a counter example with $|E(\Gamma) \cap B|$ minimal and, under that restriction, $|E(\Gamma) \setminus B|$ minimal. Obviously $|E(\Gamma) \cap B|$ is not 0 or 2. Hence $|E(\Gamma) \cap B| \geq 4$.

First we prove

If $C \in S$, $u, v \in V(C) \cap V(\Gamma)$ with $u \neq v$. Then

(*) $\min\{|E(\Pi_1) \cap B|, |E(\Pi_2) \cap B|\} = 0$, where Π_1 and Π_2 are the paths on Γ with endpoints u and v .

Assume (*) is not correct. Let P_1 and P_2 be the paths on C with endpoints u and v , such that $|E(P_1)| \equiv |E(P_2)| \pmod{2}$. Then for $i = 1, 2$, $E(P_i) \Delta E(\Pi_i)$ contains an odd cycle Γ_i , say, such that $|E(\Gamma_i) \cap B| \leq |E(\Pi_i) \cap B| < |E(\Gamma) \cap B|$. Hence $|E(P_1)| + |E(\Pi_1) \setminus B| \geq |E(\Gamma_1) \setminus B| \geq M$ (by the minimality of Γ). Since $|E(P_1)| + |E(P_2)| = M$ we now have $|E(\Gamma) \setminus B| = |E(\Pi_1) \setminus B| + |E(\Pi_2) \setminus B| \geq M$, contradicting our assumption that Γ violates claim 2, hence (*) is correct.

Now let $v_1, v_2 \in V_1 \cap V(\Gamma)$ such that the two paths Π_1 and Π_2 on Γ with end-points v_1 and v_2 satisfy $E(\Pi_1) \cap E_1 = \emptyset$, $|E(\Pi_1) \cap B| = 2$. Let C_1 and C_2 be shortest odd cycles in G_1 with $v_1 \in V(C_1)$ and $v_2 \in V(C_2)$, and such that $|E(C_i) \cap E(\Gamma)|$ is maximal for $i = 1, 2$. Then $|V(C_i) \cap V(\Gamma)| = 1$ or $E(C_i) \cap E(\Gamma)$ is a path.

Indeed, let $u \in V(C_i) \cap V(\Gamma) \setminus \{v_i\}$. Let Q be the path on Π_2 from v_i to u , and let P be the path on C_i from v_i to u , such that $|E(P)| = |E(Q)|$. Since $C_i \in S$: $|E(Q)| \geq |E(P)|$. Let Γ' be an odd cycle in $(E(\Gamma) \setminus E(Q)) \cup E(P)$. By $(*)$ $E(Q) \cap B = \emptyset$; hence $|E(\Gamma') \cap B| \leq |E(\Gamma) \cap B|$ and $|E(\Gamma') \setminus B| \leq |E(\Gamma) \setminus B| - |E(Q)| + |E(P)|$.

By the minimality of Γ we now have $|E(Q)| \leq |E(P)|$, so $|E(Q)| = |E(P)|$. Hence $(E(C_i) \setminus E(P)) \cup E(Q)$ is a shortest odd cycle, which violates the maximality of $|E(C_i) \cap E(\Gamma)|$ unless $P=Q$.

Now let $b \in E(\Pi_2) \cap E(B)$ (which is not empty). Let Π_{21} and Π_{22} be the components of $E(\Pi_2) \setminus \{b\}$ with $v_i \in V(\Pi_{2i})$, $i = 1, 2$. By $(*)$ $V(\Pi_{21}) \cap V(C_2) = \emptyset$ and $V(\Pi_{22}) \cap V(C_1) = \emptyset$. Hence $V(\Gamma) \cap V(C_1) \cap V(C_2) = \emptyset$.

Since $V(C_1) \cap V(C_2) \neq \emptyset$ (claim 1) and since $E(C_1) \cap E(\Gamma)$ is a path, the cycle Γ is non-separating in the union of Γ , C_1 and C_2 . So by the lemma there exists an odd cycle C , say, in $E(C_1) \cup E(C_2)$ such that $V(C) \cap V(\Gamma) = \emptyset$. From the fact that C_1 and C_2 are in S it is not hard to see that $C \in S$, and $(E(C_1) \cup E(C_2)) \setminus E(C)$ is also a cycle in S . This cycle violates $(*)$.

end of proof of claim 2

Now we contract the edges in B . We get a graph G' which does not contain an odd- K_4 or an odd- K_3^2 (by the proposition), and in which each odd cycle has length greater than or equal to M (claim 2). By the minimality of G , G' maps to a cycle of length M . Equivalently there exists a map $\varphi' : V(G') \rightarrow \mathbb{Z}$ such that $|\varphi'(u) - \varphi'(v)| \equiv 1 \pmod{M}$ for each $uv \in E(G')$.

Define $\varphi : V(G) \rightarrow \mathbb{Z}$ by $\varphi(u) = \varphi'(u')$ if $u \notin V_1$ and $\varphi(u) = \varphi'(u') + 1$ if $u \in V_1$, where u' is the node in G' to which u is contracted.

Claim 3. $|\varphi(u) - \varphi(v)| \equiv 1 \pmod{M}$ for each $uv \in E(G)$.

Proof of claim 3. Let $u, v \in V(G)$, $uv \in E(G)$. If $uv \in B$ then u and v are contracted into the same node of G' , and $u \in V_1$, $v \notin V_1$ (or vice versa). Hence $\varphi(u) = \varphi(v) + 1$. If $uv \notin B$ then u and v are both in V_1 or both not in V_1 . Moreover they are not contracted to one node since then there would be

an odd cycle Γ in G with $|E(\Gamma) \setminus B| = 1$, violating claim 2. So
 $|\varphi(u) - \varphi(v)| \equiv |\varphi'(u') - \varphi'(v')| \equiv 1 \pmod{M}$.

end of proof of claim 3

Hence G maps on a cycle of length M , contradicting our assumption that G violates the theorem, so our assumption $k \geq 2$ is not correct. Take any odd cycle $C \in S$. By claim 1 it meets any odd cycle in G . Since C is non-separating there exists a $v \in V(C)$, such that each odd cycle in G passes through v (see the lemma). Hence the graph induced by $V \setminus \{v\}$ is bipartite. Let $V_r \cup V_b$ be a bipartition. Denote the set of nodes of $V \setminus \{v\}$ which have distance 0 to v by V^i . The collection $\{V^i | i = 1, \dots, \lfloor \frac{1}{2}M \rfloor\}$ is a partition of $V \setminus \{v\}$ as each $u \in V$ is in an odd cycle of length M . Moreover $V^i \cap V_r$ and $V^i \cap V_b$ are stable sets. Contracting each of them to a single node yields a cycle of length M (this contraction is a homomorphism!). So again G maps to its shortest odd cycle, thus providing a final contradiction. \square

REMARK. It is not hard to deduce from the proof a polynomial-time algorithm which determines, given a graph G with no odd- K_4 or odd- K_3^2 , a homomorphic map from G to its shortest odd cycle. Indeed, the length of a shortest odd cycle, as well as a shortest odd cycle containing some fixed node or edge can be determined in polynomial-time. Hence so can E_S (Grötschel, Pulleyblank [1981] or Gerards, Schrijver [1985a]). Moreover, if we find a map to a cycle of length M after we contracted the edges leaving some component of (V, E_S) , we can easily find such a map for the original graph. In fact we can use the above proof to create a polynomial-time algorithm which determines, given a graph G , a homomorphic map from G to its shortest odd cycle, an odd- K_4 , or an odd- K_3^2 . That algorithm starts with the algorithm indicated above. If that fails (e.g. claim 2 does not hold, or E_S is connected but there is no $v \in V(G)$ such that $V(G) \setminus \{v\}$ induces a bipartite graph) then it is possible to re-construct the proof in polynomial-time, such that one finds an odd- K_4 , an odd- K_3^2 or a separating shortest odd cycle. In the last case a decomposition of the problem is possible. Since the details are rather complicated, and messy, we shall omit them.

ACKNOWLEDGEMENT

I wish to thank Alexander Schrijver, who called my attention to the subject of this paper and supported me with the preparation.

REFERENCES

- [1984] M.O. Albertson, P.A. Catlin and L. Gibbons, *Homomorphisms of 3-chromatic graphs*, II, preprint.
- [1985a] A.M.H. Gerards and A. Schrijver, *Matrices with the Edmonds-Johnson property*, Report No. 85363, Inst. für Ökon. u. Op. Res., Uni. Bonn, W. Germany.
- [1985b] A.M.H. Gerards and A. Schrijver, in preparation.
- [1981] M. Grötschel and W.R. Pulleyblank, *Weakly bipartite graphs and the max-cut problem*, Oper. Res. Letters 1 (1981), 23-27.
- [1977] P.D. Seymour, *The matroids with the max-flow min-cut property*, J. Combinatorial theory (B) 23 (1977) 189-222.
- [1958] W.T. Tutte, *A homotopy theorem for matroids*, II, Trans. Amer. Math. Soc., 88 (1958) 161-174.

IN 1984 REEDS VERSCHENEN

- 138 G.J. Cuypers, J.P.C. Kleijnen en J.W.M. van Rooyen
Testing the Mean of an Asymmetric Population:
Four Procedures Evaluated
- 139 T. Wansbeek en A. Kapteyn
Estimation in a linear model with serially correlated errors when
observations are missing
- 140 A. Kapteyn, S. van de Geer, H. van de Stadt, T. Wansbeek
Interdependent preferences: an econometric analysis
- 141 W.J.H. van Groenendaal
Discrete and continuous univariate modelling
- 142 J.P.C. Kleijnen, P. Cremers, F. van Belle
The power of weighted and ordinary least squares with estimated
unequal variances in experimental design
- 143 J.P.C. Kleijnen
Superefficient estimation of power functions in simulation
experiments
- 144 P.A. Bekker, D.S.G. Pollock
Identification of linear stochastic models with covariance
restrictions.
- 145 Max D. Merbis, Aart J. de Zeeuw
From structural form to state-space form
- 146 T.M. Doup and A.J.J. Talman
A new variable dimension simplicial algorithm to find equilibria on
the product space of unit simplices.
- 147 G. van der Laan, A.J.J. Talman and L. Van der Heyden
Variable dimension algorithms for unproper labellings.
- 148 G.J.C.Th. van Schijndel
Dynamic firm behaviour and financial leverage clienteles
- 149 M. Plattel, J. Peil
The ethico-political and theoretical reconstruction of contemporary
economic doctrines
- 150 F.J.A.M. Hoes, C.W. Vroom
Japanese Business Policy: The Cash Flow Triangle
an exercise in sociological demystification
- 151 T.M. Doup, G. van der Laan and A.J.J. Talman
The $(2^{n+1}-2)$ -ray algorithm: a new simplicial algorithm to compute
economic equilibria

IN 1984 REEDS VERSCHENEN (vervolg)

- 152 A.L. Hempenius, P.G.H. Mulder
Total Mortality Analysis of the Rotterdam Sample of the Kaunas-Rotterdam Intervention Study (KRIS)
- 153 A. Kapteyn, P. Kooreman
A disaggregated analysis of the allocation of time within the household.
- 154 T. Wansbeek, A. Kapteyn
Statistically and Computationally Efficient Estimation of the Gravity Model.
- 155 P.F.P.M. Nederstigt
Over de kosten per ziekenhuisopname en levensduurmodellen
- 156 B.R. Meijboom
An input-output like corporate model including multiple technologies and make-or-buy decisions
- 157 P. Kooreman, A. Kapteyn
Estimation of Rationed and Unrationed Household Labor Supply Functions Using Flexible Functional Forms
- 158 R. Heuts, J. van Lieshout
An implementation of an inventory model with stochastic lead time
- 159 P.A. Bekker
Comment on: Identification in the Linear Errors in Variables Model
- 160 P. Meys
Functies en vormen van de burgerlijke staat
Over parlementarisme, corporatisme en autoritair etatisme
- 161 J.P.C. Kleijnen, H.M.M.T. Denis, R.M.G. Kerckhoffs
Efficient estimation of power functions
- 162 H.L. Theuns
The emergence of research on third world tourism: 1945 to 1970;
An introductory essay cum bibliography
- 163 F. Boekema, L. Verhoef
De "Grijze" sector zwart op wit
Werklozenprojecten en ondersteunende instanties in Nederland in kaart gebracht
- 164 G. van der Laan, A.J.J. Talman, L. Van der Heyden
Shortest paths for simplicial algorithms
- 165 J.H.F. Schilderink
Interregional structure of the European Community
Part II: Interregional input-output tables of the European Community 1959, 1965, 1970 and 1975.

IN (1984) REEDS VERSCHENEN (vervolg)

- 166 P.J.F.G. Meulendijks
An exercise in welfare economics (I)
- 167 L. Elsner, M.H.C. Paardekooper
On measures of nonnormality of matrices.

IN 1985 REEDS VERSCHENEN

- 168 T.M. Doup, A.J.J. Talman
A continuous deformation algorithm on the product space of unit
simplices
- 169 P.A. Bekker
A note on the identification of restricted factor loading matrices
- 170 J.H.M. Donders, A.M. van Nunen
Economische politiek in een twee-sectoren-model
- 171 L.H.M. Bosch, W.A.M. de Lange
Shift work in health care
- 172 B.B. van der Genugten
Asymptotic Normality of Least Squares Estimators in Autoregressive
Linear Regression Models
- 173 R.J. de Groof
Geïsoleerde versus gecoördineerde economische politiek in een twee-
regiomodel
- 174 G. van der Laan, A.J.J. Talman
Adjustment processes for finding economic equilibria
- 175 B.R. Meijboom
Horizontal mixed decomposition
- 176 F. van der Ploeg, A.J. de Zeeuw
Non-cooperative strategies for dynamic policy games and the problem
of time inconsistency: a comment
- 177 B.R. Meijboom
A two-level planning procedure with respect to make-or-buy deci-
sions, including cost allocations
- 178 N.J. de Beer
Voorspelprestaties van het Centraal Planbureau in de periode 1953
t/m 1980
- 178a N.J. de Beer
BIJLAGEN bij Voorspelprestaties van het Centraal Planbureau in de
periode 1953 t/m 1980
- 179 R.J.M. Alessie, A. Kapteyn, W.H.J. de Freytas
De invloed van demografische factoren en inkomen op consumptieve
uitgaven
- 180 P. Kooreman, A. Kapteyn
Estimation of a game theoretic model of household labor supply
- 181 A.J. de Zeeuw, A.C. Meijdam
On Expectations, Information and Dynamic Game Equilibria

- 182 Cristina Pennavaja
Periodization approaches of capitalist development.
A critical survey
- 183 J.P.C. Kleijnen, G.L.J. Kloppenburg and F.L. Meeuwsen
Testing the mean of an asymmetric population: Johnson's modified T
test revisited
- 184 M.O. Nijkamp, A.M. van Nunen
Freia versus Vintaf, een analyse

Bibliotheek K. U. Brabant



17 000 01059755 8